

The more you know, the less sure you are.

- Voltaire

1. [University of Michigan] For each of the following three sets of axes, exactly one of the following statements **(A)-(E)** is true. You may use a letter more than once. In the space provided next to each figure, enter the letter of the true statement for that figure. For each graph, note that the x and y scales are not the same.

(A) h is the derivative of f , and f is the derivative of g .


(B) g is the derivative of f , and f is the derivative of h .

(C) g is the derivative of h , and h is the derivative of f .

(D) h is the derivative of g , and g is the derivative of f .


(E) none of (A) through (D) are possible

(G1)




ANSWER: A

(G2)



ANSWER: E

(G3)



ANSWER: D

2. Consider the rational function $f(x) = \frac{x(3x+1)^2(2x+3)^4}{(x-5)^7}$

(a) Find the *domain* of f .

Solution: $f(x)$ is defined for all real numbers except 5

(b) Find the y -intercept of f .

Solution: The y -intercept of f is $f(0) = 0$.

(c) Find the zeroes of f .

Solution: $f(x) = 0$ when $x = 0, -1/3, -3/2$

(d) Find the vertical asymptote(s) of f .

Solution: A unique singularity occurs at $x = 5$; hence the only vertical asymptote is the line $x = 5$.

(e) Find the horizontal asymptote(s) of f .

Solution: For large $|x|$,

$$f(x) = \frac{x(3x+1)^2(2x+3)^4}{(x-5)^7} \approx \frac{x(3x)^2(2x)^4}{(x)^7} = \frac{x(9x^2)(16x)^4}{x^7} = 144$$

Thus the only horizontal asymptote is the line: $y = 144$

3. Albertine orders a large cup of coffee at Metropolis on Granville. Let $F(t)$ be the temperature in *degrees Fahrenheit* of her coffee t minutes after the coffee is placed on her tray.

(a) Explain the meaning of the statement: $F(9) = 167$. (Use complete sentences. Avoid any mathematical terms!)

Solution: Nine minutes after being served her coffee, Albertine finds that the coffee is 167°F .

(b) Explain the meaning of the statement: $F^{-1}(99) = 17.5$

Solution: When the temperature of the coffee is 99°F , 17.5 minutes have elapsed since the coffee mug was placed on her tray.

(c) Give the *practical* interpretation of the statement: $F'(9) = -1.10$. (Use complete sentences. Do not use the words “derivative” or “rate” or any other mathematical term in your explanation.)

Solution: Nine minutes after being served her coffee, Albertine notices that the temperature of the coffee is decreasing by about 0.11°F every tenth of a minute during the next minute or two.

(d) What are the *units* of $F'(9)$?

*Solution: $^{\circ}\text{F}/\text{minute}$ (since $\Delta F/\Delta t$ represents *change in temp/ change in time*.)*

(e) Using the information given in parts (a) and (c), estimate the temperature of Albertine’s coffee seven minutes after she has been handed the coffee.

Solution: Since after nine minutes the temperature is 167°F , we would estimate that two minutes earlier the temperature was $167 + 2(1.1) = 169.2^{\circ}\text{F}$.

(e) **EXTRA CREDIT**

Explain the meaning of the statement:

$$(F^{-1})'(99) = -1$$

Solution: When the temperature of the coffee is 99°F , the time required for the temperature of the coffee to fall to 98°F is one additional minute.

4. The following function is *not* continuous at the points $x = 5/2$ and $x = 4/3$.

$$g(x) = \frac{2x^5 - 19x^4 + 35x^3}{6x^2 - 23x + 20}$$

- (a) [6 pts] Does g possess a *continuous extension* at $x = 5/2$? If so, how should $g(5/2)$ be defined? If not, explain!

Solution:

We begin by factoring:

$$g(x) = \frac{2x^5 - 19x^4 + 35x^3}{6x^2 - 23x + 20} = \frac{x^3(2x - 5)(x - 7)}{(3x - 4)(2x - 5)}$$

If $x \neq 5/2$, then cancellation yields:

$$g(x) = \frac{x^3(x - 7)}{3x - 4}$$

So we see that $x = 5/2$ is a removable discontinuity.

To create a continuous extension to g at $x = 5/2$, we define $g(5/2)$ as follow:

$$g(5/2) = \frac{(5/2)^3(5/2 - 7)}{3(5/2) - 4} = -\frac{1125}{56} \cong -20.09$$

- (b) [6 pts] Does g possess a *continuous extension* at $x = 4/3$? If so, how should $g(4/3)$ be defined? If not, explain!

Solution:

We begin by noting that, when $x \neq 5/2$,

$$g(x) = \frac{x^3(x - 7)}{3x - 4}$$

Now, as $x \rightarrow 4/3$, the numerator tends toward $(4/3)^3(4/3 - 7) \cong -13.43$, but the denominator tends toward 0. Consequently, the limit of $g(x)$ as $x \rightarrow 4/3$ does not exist. (This is an infinite discontinuity.)

5. [University of Michigan] Suppose $f(x) = \left(x + \frac{1}{2}\right)e^x$

- a. Using the limit definition of the derivative, write an explicit expression for $f'(2)$. Your expression should not contain the letter “ f ”. Do not try to evaluate your expression.

$$\text{Solution: } f'(2) = \lim_{h \rightarrow 0} \frac{(2+h+\frac{1}{2})e^{2+h} - (2+\frac{1}{2})e^2}{h}$$

- b. Albertine discovers that the derivative of $f(x)$ is $f'(x) = \left(x + \frac{3}{2}\right)e^x$.

Using this formula for $f'(x)$, write an equation for the tangent line to the graph of $f(x)$ at $x = 2$.

Solution: The slope is $m = (2 + \frac{3}{2})e^2 = \frac{7}{2}e^2$ and the y-coordinate when $x = 2$ is $f(2) = (2 + \frac{1}{2})e^2 = \frac{5}{2}e^2$. Using the point-slope form for the tangent line, we obtain:

$$y - \frac{5e^2}{2} = \frac{7}{2}e^2(x - 2)$$

- c. [3 points] Write an equation for the tangent line to the graph of $f(x)$ at $x = a$ where a is an unknown constant.

Solution: Employing virtually the same computation that we did in (b), we find:

$$y - \left(a + \frac{1}{2}\right)e^a = \left(a + \frac{3}{2}\right)e^a(x - a)$$

- d. **EXTRA CREDIT** Using your answer from (c), find a value of a so that the tangent line to the graph of $f(x)$ at $x = a$ passes through the origin.

Solution: The point $(0, 0)$ must satisfy the tangent line equation in (e). Hence

$$0 - \left(a + \frac{1}{2}\right)e^a = \left(a + \frac{3}{2}\right)e^a(0 - a)$$

Dividing each side by e^a we obtain:

$$-\left(a + \frac{1}{2}\right) = -a\left(a + \frac{3}{2}\right)$$

So $-2a - 1 = -2a^2 - 3a$

Simplifying: $2a^2 + a - 1 = 0$

Factoring: $(2a - 1)(a + 1) = 0$

Hence $a = 1/2$ or $a = -1$

6. Compute each of the following limits. *Justify your reasoning.*

(a) $\lim_{x \rightarrow 1} \frac{(x-2)^5}{x^{2015} + 4x - 6}$

Solution: Using the laws of limits:

$$\lim_{x \rightarrow 1} \frac{(x-2)^5}{x^{2015} + 4x - 6} = \frac{\lim_{x \rightarrow 1} (x-2)^5}{\lim_{x \rightarrow 1} (x^{2015} + 4x - 6)} = \frac{(-1)^5}{1 + 4 - 6} = \frac{-1}{-1} = 1$$

(b) $\lim_{x \rightarrow 2} \left(1 + \frac{8}{x} + \frac{x}{2} - 5 \sin(\pi x) + 7 \cos(\pi x)\right)$

Solution: Using the laws of limits:

$$\lim_{x \rightarrow 2} \left(1 + \frac{8}{x} + \frac{x}{2} - 5 \sin(\pi x) + 7 \cos(\pi x)\right) =$$

$$\lim_{x \rightarrow 2} 1 + \lim_{x \rightarrow 2} \frac{8}{x} + \lim_{x \rightarrow 2} \frac{x}{2} - \lim_{x \rightarrow 2} 5 \sin(\pi x) + \lim_{x \rightarrow 2} 7 \cos(\pi x) =$$

$$1 + \frac{8}{2} + \frac{2}{2} - 5 \sin(2\pi) + 7 \cos(2\pi) = 1 + 4 + 1 - 5(0) + 7(1) = 13$$

$$(c) \lim_{x \rightarrow \infty} \frac{x^4(3x-1)(5x^2-19x+1)^2}{(x-2)^7(4x-13)(x+43)}$$

Solution: For large values of x :

$$\frac{x^4(3x-1)(5x^2-19x+1)^2}{(x-2)^7(4x-13)(x+43)} \approx \frac{x^4(3x)(5x^2)^2}{(x)^7(4x)(x)} = \frac{75x^9}{4x^9} = \frac{75}{4}$$

$$(d) \lim_{x \rightarrow 0} \frac{\sin(5x)}{\tan(9x)}$$

Solution:

$$\frac{\sin(5x)}{\tan(9x)} = \frac{\sin(5x)}{\sin(9x)} \cos(9x) = \frac{5}{9} \left(\frac{\frac{\sin(5x)}{5x}}{\frac{\sin(9x)}{9x}} \right) \cos(9x) \rightarrow \frac{5}{9} \left(\frac{1}{1} \right) 1 = \frac{5}{9} \text{ as } x \rightarrow 0$$

7. (a) [6 pts] Find an equation of the *normal line* to the curve

$$y = g(x) = \frac{x^2-1}{x^2+1} \text{ at } x = 1.$$

Solution:

$$dy/dx = \frac{(x^2+1)d/dx(x^2-1) - (x^2-1)d/dx(x^2+1)}{(x^2+1)^2} =$$

$$\frac{(x^2+1)(2x) - (x^2-1)(2x)}{(x^2+1)^2}$$

$$\text{Now, when } x = 1, dy/dx = \frac{(1^2+1)(2) - (1^2-1)(2)}{(1^2+1)^2} = 1$$

Hence the slope of the normal line is $m = -1$.

Also, note that $g(1) = 0$.

Using the point-slope form of the line: $y - 0 = -1(x - 1)$

or $y = 1 - x$

(b) [6 pts] Find an equation of the *tangent line* to the curve $y = F(x) = e^x (x^2 + 4x + 1)$ at $x = 0$.

Solution:

$$\begin{aligned} dy/dx &= e^x d/dx(x^2 + 4x + 1) + (x^2 + 4x + 1) d/dx(e^x) = \\ &e^x(2x + 4) + (x^2 + 4x + 1)e^x \end{aligned}$$

At $x = 0$, $dy/dx = e^0(4) + (1)e^0 = 5$.

Also note that $f(0) = e^0 = 1$.

So the equation of the tangent line at the point $(0, 1)$ is given by:


$$y - 1 = 5(x - 0)$$

or $y = 5x + 1$

8. Identify the *type of discontinuity* that each of the following functions has at $x = 0$. (Choose from: *removable*, *infinite*, *jump*, or *essential* discontinuity.) You need not justify your answers.


$$(a) \quad y = \frac{x^5 - x^4 + x^3 - x^2 + x}{x^2 + x}$$

removable discontinuity at $x = 0$




$$(b) \quad y = \cos\left(\frac{\sin x}{x}\right)$$

removable discontinuity at $x = 0$




$$(c) \quad y = \sinh\left(\frac{1}{x}\right)$$

infinite discontinuity at $x = 0$



(d) $y = e^{-1/x^2}$

removable discontinuity at $x = 0$



9. (a) Let $G(x) = a e^{bx}$, where a and b are non-zero constants. Albertine, our friend, informs us that $(d/dx) e^{bx} = b e^{bx}$. Find $G^{(2015)}(x)$.

Solution:

$$G'(x) = ab e^{bx}$$

$$G''(x) = (d/dx) G'(x) = (d/dx) (ab e^{bx}) = ab^2 e^{bx}$$

$$G'''(x) = (d/dx) G''(x) = (d/dx) (ab^2 e^{bx}) = ab^3 e^{bx}$$

$$G^{(4)}(x) = ab^4 e^{bx}$$

$$G^{(5)}(x) = ab^5 e^{bx}$$


Based upon this pattern, we infer that $G^{(n)}(x) = ab^n e^{bx}$

$$\text{Thus } G^{(2015)}(x) = ab^{2015} e^{2015bx}$$

(b) [University of Michigan] Your pet parakeet, Odette, is flying in a straight path that at first is toward you and then away from you for a minute. After t seconds, she is $f(t)$ feet away from you, where

$$f(t) = \frac{-t(t-20)(t-70)}{500} + 20, \quad 0 \leq t \leq 60.$$

A graph of $y = f(t)$ is shown below.




Without doing any calculations, determine which is greater: the *average velocity of Odette over the entire minute*, or her *instantaneous velocity at 30 seconds*. Explain, referring to the graph. (The absence of a clear explanation will result in no credit.)

Solution:

The slope of the secant line from $t = 0$ to $t = 60$ is the average velocity over the minute and the slope of the tangent line at $t = 30$ is the instantaneous velocity after 30 seconds. If you draw these lines on the graph, you can see that the tangent line clearly has a larger slope. Thus, Odette's instantaneous velocity after 30 seconds is greater than her average velocity over the entire minute.

10. (a) State carefully the *Intermediate Value Theorem* for a function $y = G(x)$. Provide a sketch that illustrates the statement.

Let $y = f(x)$ be a continuous function on the interval $[a, b]$. Let u be any number between $f(a)$ and $f(b)$. Then there exists a number c in the interval $[a, b]$ such that $f(c) = u$.



(b) Let $G(x) = x^5 - 5 - \ln x$. Using the *Intermediate Value Theorem*, carefully explain why the equation $G(x) = 0$ has *at least one* solution. (Your calculator is not needed here.)

Solution:

Note that $G(1) = 1 - 5 - \ln 1 = -4 < 0$ and $G(e) = e^5 - 5 - \ln e = e^5 - 6 > 2^5 - 6 = 28 > 0$.

*Also note that G is continuous on the interval $[0, \pi]$. Since $G(0) < 0 < G(e)$, the *IVT* guarantees the existence of a root of the equation $G(x) = 0$ in the interval $[0, e]$.*

11. Suppose that f and g are differentiable functions satisfying:

$$f(3) = -2, g(3) = -4, f'(3) = 3, \text{ and } g'(3) = -1.$$

(a) Let $H(x) = (f(x) + 2g(x) + 1)(f(x) - g(x) - 4)$. Compute $H'(3)$

Solution: Using the product rule:

$$\begin{aligned} H'(x) &= (f(x) + 2g(x) + 1) \frac{d}{dx}(f(x) - g(x) - 4) + (f(x) - g(x) - 4) \frac{d}{dx}(f(x) + 2g(x) + 1) \\ &= (f(x) + 2g(x) + 1) (f'(x) - g'(x)) + (f(x) - g(x) - 4)(f'(x) + 2g'(x)) \end{aligned}$$

$$\begin{aligned} \text{Thus } H'(3) &= (f(3) + 2g(3) + 1) (f'(3) - g'(3)) + (f(3) - g(3) - 4)(f'(3) + 2g'(3)) \\ &= (-2 - 8 + 1) (3 - (-1)) + (-2 - (-4) - 4)(3 + 2(-1)) = (-9)(4) + (-2)(1) = -38 \end{aligned}$$

(b) Let $M(x) = \frac{2f(x) + 3g(x)}{2 - 3g(x)}$. Compute $M'(3)$

Solution: Using the quotient rule:


$$M'(x) = \frac{(2 - 3g(x)) \frac{d}{dx}(2f(x) + 3g(x)) - (2f(x) + 3g(x)) \frac{d}{dx}(2 - 3g(x))}{(2 - 3g(x))^2} =$$

$$\frac{(2 - 3g(x))(2f'(x) + 3g'(x)) - (2f(x) + 3g(x))(-3g'(x))}{(2 - 3g(x))^2}$$

At $x = 3$, $M'(3) =$

$$\frac{(2 - 3(-4))(2(3) + 3(-1)) - (2(-2) + 3(-4))(-3(-1))}{(2 - 3(-4))^2} = \frac{45}{98}$$

12. [University of Michigan] The graph of a function $h(x)$ is given below.



- a. [2 point] List all x -values with $-4 < x < 4$ where $h(x)$ is not continuous. If there are none, write NONE.

Solution: $h(x)$ is not continuous at $x = 1$. Notice it is a jump discontinuity.

- b. [2 point] List all x -values with $-4 < x < 4$ where $h(x)$ is not differentiable. If there are none, write NONE.

Solution: $h(x)$ is not differentiable at two points: $x = 1$ and $x = 3$.

- c. On the axes provided, carefully draw a graph of $y = h'(x)$. Be sure to label important points or features on your graph.

