

# MATH 162

# PRACTICE TEST II

1. For each of the following infinite series, determine *convergence* or *divergence*. In the case of convergence, find the sum of the series:

$$(a) \sum_{n=1}^{\infty} \ln \frac{n+1}{n}$$

$$(b) \sum_{n=0}^{\infty} \frac{5}{9^n}$$

$$(c) \sum_{n=1}^{\infty} \left( \frac{n}{n+1} \right)^n$$

$$(d) \sum_{n=1}^{\infty} \frac{1}{n(n+1)}$$

$$(e) \sum_{n=1}^{\infty} \cos \left( \frac{5}{n} \right)$$

$$(f) 0.123123123\dots$$

2. For each series below, determine absolute convergence, conditional convergence or divergence. Justify each answer.

$$(a) \sum_{n=3}^{\infty} (-1)^n \frac{13}{(\ln n)^{13}}$$

$$(b) \sum_{k=1}^{\infty} (-1)^k \frac{(k+3)(k^2+5)}{(k+13 \ln k)^4}$$

$$(c) \sum_{n=1}^{\infty} (-1)^n \frac{2^n}{\left(1 + \frac{1}{n}\right)^{n^2}}$$

$$(d) \sum_{n=1}^{\infty} (-1)^n \frac{1}{\ln(e^n + e^{-n})}$$

$$(e) \sum_{n=1}^{\infty} (-1)^n \frac{n^{13}}{(n+13)!}$$

3. For each power series below, determine the *radius of convergence* and the *interval of convergence*. Study the behavior of each power series at the *endpoints*.

$$(a) \sum_{n=1}^{\infty} \frac{13^n}{n(n+13)} x^n$$

$$(b) \sum_{n=1}^{\infty} \frac{1}{n(n+3)(n+11)} (x-4)^n$$

$$(c) \sum_{n=1}^{\infty} \frac{1}{\sqrt{3n+7}} (x+13)^n$$

4. (a) Find the 3<sup>rd</sup> order Maclaurin polynomial of  $\cosh x$ .

(b) Find the 5<sup>th</sup> order Taylor polynomial of  $\cos x$  centered at  $x = \pi/2$ .

5. Find the 4<sup>th</sup> order Taylor polynomial of  $e^x$  centered at  $x = 2$ .

6. Find the 3<sup>rd</sup> order Maclaurin polynomial of

$$f(x) = 4 + (x+13)^2 + (x+13)^3$$

7. By differentiating the power series expansion of  $y = 1/(1-x)$ , find the value of

$$\sum_{k=0}^{\infty} \frac{k}{13^k}$$

8. Find the *first five* non-zero terms of the Maclaurin series expansion of

$$h(x) = (1 + 2x^2) e^{3x}.$$

9. Let  $f(x) = x^8 e^{5x}$ . Compute  $f^{(100)}(0)$ . Do not simplify your answer.

10. Find the *radius of convergence* of the power series:

$$\sum_{n=1}^{\infty} \frac{n!}{(1)(3)(5)(7)\dots(2n-1)} x^n$$

11. Find the *radius of convergence* of the power series:

$$\sum_{n=0}^{\infty} n! x^{2^n}$$

12. Find the *radius of convergence* of the power series:

$$\sum_{n=1}^{\infty} \frac{1}{(\ln n)^n} x^n$$

13. Without using l'Hôpital's rule, calculate the following limit. Show your work!

$$\lim_{t \rightarrow 0} \frac{te^{4t} - \sin(3t) + 2t - 4t^2}{t^3}$$

14. Let  $G(x) = x^3 \cosh(3x)$ . Using an appropriate Maclaurin series, compute  $G^{(2017)}(0)$ . (Do not try to simplify your answer.)

15. Find the first four non-zero terms of the Maclaurin series for each of the following:

(a)  $\frac{e^{2x}}{\cosh x}$

(b)  $\frac{\ln(1+x)}{1+x^2}$

(c)  $e^{x^2} \sin 2x$

16. Using division of power series, find the first three non-zero terms of the Maclaurin series expansion of

$$f(x) = \frac{e^{2x} + 1}{\cos x}$$

17. Using multiplication of power series, find the first four non-zero terms of the Maclaurin series expansion of

$$g(x) = e^{x^2} (1 + x^2 + x^3)$$

18. Determine the *interval of convergence* of the following power series. (You need not study end-point behavior.)


$$\sum_{n=1}^{\infty} \frac{n^{13} 13^n}{\sqrt{n+2016}} (x-13)^n$$

19. Analyze the behavior of the series

$$\sum_{n=1}^{\infty} \frac{\sqrt{n^2 + 4}}{(n^{1/3} + 1789)^5}$$

20. What is the relationship between  $\cosh x$  and  $\cos x$ ? between  $\sinh x$  and  $\sin x$ ? State Euler's formula.

21. [University of Michigan final exam question] The graph shows the area between the graphs of  $f(x) = 6 \cos(\sqrt{2}x)$  and  $g(x) = x^2 + x$ . Let  $(x_0, y_0)$  be the intersection point between the graphs of  $f(x)$  and  $g(x)$ .



(a) Compute  $P(x)$ , the function containing the first three nonzero terms of the Taylor series about  $x = 0$  of  $f(x) = 6 \cos(\sqrt{2}x)$ .

(b) Use  $P(x)$  to approximate the value of  $x_0$ .

(c) Use  $P(x)$  and the value of  $x_0$  you computed in the previous question to write an integral that approximates the value of the shaded area.

(d) Graph  $f(x)$  and  $g(x)$  in Wolfram Alpha or your calculator. Use the graphs to find an approximate value for  $x_0$ .

(e) Write a definite integral in terms of  $f(x)$  and  $g(x)$  that represents the value of the shaded area. Find its value using your Wolfram Alpha or your calculator.

22. [University of Michigan final exam question] (a) Find the Maclaurin series of  $\sin(x^2)$ . Your answer should include a formula for the general term in the series.

(b) Let  $m$  be a positive integer, find the Maclaurin series of  $\cos(m\pi x)$ . Your answer should include a formula for the general term in the series.

(c) Use the second degree Maclaurin polynomials of  $\sin(x^2)$  and  $\cos(m\pi x)$  to approximate the value of  $b_m$ , where

$$b_m = \int_{-1}^1 \sin(x^2) \cos(m\pi x) dx.$$

(The number  $b_m$  is called a *Fourier coefficient of the function*  $\sin(x^2)$ . These numbers play a key role in Fourier analysis, a subject with widespread applications in engineering and the sciences.)



Jean-Baptiste Joseph Fourier (1768 – 1830)