MATH 201: PREPARING FOR TEST 2 (REVISED 3/11/2019)

Chapter 3 (sections 1 – 5, 7 (special case), 8 – 10); Chapter 10 (section 1); Chapters 4 – 6; Chapter 7 (as much as we cover Monday & Wednesday)



Albrecht-Dürer, Melancholia

(Note the magic square in the background.)

- > Types of problems:
 - counting: permutations, combinations, stars & bars
 - proofs (direct, cases, induction (ordinary), contrapositive, contradiction, if and only if, conditional)
 - definitions in the case of number theory (odd, even, divisibility, prime number, etc.)
 - statements of theorems (pigeon-hole principle, inclusion/exclusion, division algorithm, etc.)
 - fill in the blank
 - True/False
 - find a counter-example
 - given proof with a missing part, fill in the missing part
 - given a false "proof," correct it
 - modular arithmetic
 - pigeon-hole principle
 - inclusion/exclusion principle (special case)

Practice problems:

- 1. Consider the set of all binary sequences of length 10. How many such strings exist if
 - (a) No condition?
 - (b) Exactly three 1s?
 - (c) First digit and last digit must be 0?
 - (d) First digit or last digit must be 0?
 - (e) The sum of the digits is 7?
- **2.** Let n be an integer. Prove that $n^3 + n^2 + n$ is even if and only if n is even.
- 3. Given integers, p and q prove that if both pq and p+q are even, then both p and q are even using
 - (a) Proof by contrapositive
 - (b) Direct proof
- 4. Given integers c and d, where $c \ge 2$, prove, using the method of contradiction, that either $c \nmid d$ or $c \nmid (d + 1)$.
- 5. Let p and q be integers and let m be an integer greater than or equal to 1. Then we write $p \equiv q \pmod{m}$ if

6. The basic version of the pigeonhole principle states that if there are n-pigeon holes, k pigeons and ______ then

7. Let a and b be non-zero integers. Then we say write a|b if ______.

8. Let n and d be integers with $d \neq 0$. The *division algorithm* states that there exist unique integers q and r such that

______ where ______.

- **9.** The two basic steps in a proof by induction are called:
 - (A) _____

- (B) _____
- **10.** 15⁹⁹⁹⁹⁹ (mod 16) is congruent to _____
- **11.** If $a \equiv 9 \pmod{11}$ and $b \equiv 7 \pmod{11}$, then $3a + 5b \equiv (\mod{11})$
- 12. Explain why every integer can be expressed in the form 5n, 5n+1, 5n+2, 5n+3 or 5n+4.
- **13.** Find the units digit of 17^{902} . (*Hint:* Think mod 10.)
- **14.** There are 800,000 pine trees in a forest. Each pine tree has no more than 600,000 needles. Prove that at least two trees have the same number of needles.
- 15. Harvey Swick Middle School has 1,837 pupils currently registered.Prove that at least 6 of the students celebrate their birthdays on the same day of the year.

16. Let a, b, c, d and m be integers, with $m \ge 2$. Prove that if $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$

- **17.** Find 1! + 2! + 3! + ... + 12345! (mod 12)
- **18.** Let a and c be *positive* integers and let $m \ge 2$ be an integer. Demonstrate, providing a counter-example, that if $ca \equiv cb \pmod{m}$ then it needn't follow that $a \equiv b \pmod{m}$.
- **19.** Let a, b and c be positive integers. Prove that if ca|cb then a|b.

20. Find the flaw in the following "proof":

(from A. J. Hildebrand, notes from University of Illinois)

Claim: For all $n \in \mathbb{N}$, $(*) \sum_{i=1}^{n} i = \frac{1}{2}(n + \frac{1}{2})^2$

Proof: We prove the claim by induction.

Base step: When n = 1, (*) holds.

Induction step: Let $k \in \mathbb{N}$ and suppose (*) holds for n = k. Then

$$\sum_{i=1}^{k+1} i = \sum_{i=1}^{k} i + (k+1)$$

= $\frac{1}{2} \left(k + \frac{1}{2} \right)^2 + (k+1)$ (by ind. hypothesis)
= $\frac{1}{2} \left(k^2 + k + \frac{1}{4} + 2k + 2 \right)$ (by algebra)
= $\frac{1}{2} \left(\left(k + 1 + \frac{1}{2} \right)^2 - 3k - \frac{9}{4} + k + \frac{1}{4} + 2k + 2 \right)$ (more algebra)
= $\frac{1}{2} \left((k+1) + \frac{1}{2} \right)^2$ (simplifying).

Thus, (*) holds for n = k + 1, so the induction step is complete.

Conclusion: By the principle of induction, (*) holds for all $n \in \mathbb{N}$.

- **21.** Prove that $111^{333} + 333^{111}$ is divisible by 7.
- **22.** Fifteen students in French 103 were given a dictation quiz. Albertine made 13 errors. Each of the other students made fewer errors. Prove that *at least two* students made the same number of errors. (*Who are the pigeons and what are the pigeon holes?*)
- **23.** In class, we proved that, for all natural numbers *n*, any $2^n \times 2^n$ punctured chessboard can be tiled by tri-ominos. Prove independently that 3 must be a divisor of $2^{2n} - 1$
- 24. Using modular arithmetic, find the remainder when
 - (a) 2^{125} is divided by 7.
 - (b) (12)(29)(408) is divided by 13
 - (c) 7^{1942} is divided by 5.

Restate each of the above as a statement in modular arithmetic.

25. (a) If it is now 2:00, what time would it be in 12345 hours?

- (b) Is $2222^{5555} + 5555^{2222}$ divisible by 7?
- **26.** Albertine is teaching Chem 105 this semester. She has a total of 100 students in her class. Taking a survey (via Piazza) she finds that 28 students like dogs, 26 like cats, and 16 like rats. There are 12 students who like cats and dogs, 4 who like dogs and rats, and 6 who like cats and rats. Only 2 students like dogs, cats and rats. How many students do not like any of the 3 animals: dogs, cats, rats.





Three distinguishable dice are thrown. In how many ways can the maximum of the 3

numbers occurring equal 5?

27.

- 28. If you are dealt a hand of 7 cards from a standard deck (without regard to order), how many ways can you have: (a) A flush?
 - (b) 3 pairs (excluding 4 of a kind)?
 - (c) 4 of a kind and one pair (excluding 3 of a kind)?
 - (d) 4 of a kind and no other pair?
 - (e) No pair at all?

29. Prove that $2^{1/3}$ is irrational.



Gambling was in his blood. Couldn't help it – he had an irregular heartbeat. Even his pulse was impulsive.